



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 63/00, 25/26, 25/22	A1	(11) International Publication Number: WO 98/15183 (43) International Publication Date: 16 April 1998 (16.04.98)
(21) International Application Number: PCT/US97/17393 (22) International Filing Date: 29 September 1997 (29.09.97) (30) Priority Data: 60/027,512 7 October 1996 (07.10.96) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): FOSTER, James, Perry [US/US]; 9 Cook Court, Avondale, PA 19311 (US). (74) Agent: BIRCH, Linda, D.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: A PROCESS FOR COATING BIOLOGICAL PESTICIDES AND COMPOSITIONS THEREFROM (57) Abstract A pesticidal composition comprising particles of biological pesticides and particles of durable TiO ₂ in an amount sufficient to substantially coat the particles of biological pesticides and process for the preparation thereof are disclosed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

A PROCESS FOR COATING BIOLOGICAL PESTICIDES AND COMPOSITIONS
THEREFROM

FIELD OF THE INVENTION

5 The present invention relates to biological pesticide compositions comprising particles of biological pesticides coated with durable TiO_2 and advantageous processes useful for preparing them.

BACKGROUND OF THE INVENTION

10 One aspect of the present invention relates to baculoviruses which are known to have insecticidal activity. Their commercial utility has been limited though because, without special formulation, they are readily deactivated by the UV radiation in sunlight. The incorporation of UV protectants in a variety of ways is known and some can enhance the stability of these viruses somewhat, however such compositions can be complicated, expensive to prepare and further improvements in stability and
15 compatibility with the virus are needed.

 One method of stabilizing the virus is to mix a water soluble UV absorbing substance with the virus in a spray tank. One such material is a sulfonated copolymer of catechin and leucocyanidin (U.S. Patent 4,094,969). Although some degree of
20 stabilization is achieved, there is no bond between the virus and the protecting substance hence rainfall or dew may readily separate them. Most simple tank mixes of ingredients suffer from this general problem and do not offer sufficient crop protection due to the lack of UV stability of these viruses.

 Microencapsulation using a variety of techniques is another method used. However these systems are limited due to the virus's sensitivity to many encapsulating
25 precursors and the overall expense of the preparations. Many, owing to their transparency in the UV-A and -B region of sunshine, require an additional light screening agent. Typically the screening agent is incorporated into the capsule wall such as in U.S. Patents 4,844,896 and 4,948,586; EP 0 653 158 A1; and WO 96/03041. A major problem with these preparations is the lack of a suitable triggered release
30 matrix. In other words, the encapsulated virus is constrained in a matrix which requires some process to occur for it to be released at the site of action. These release processes typically include hydrolysis, desorption or dispersion and by their nature take time to occur. Due to the limited residence time of the virus in the insect's midgut, the site of infection, fast and complete release is necessary to reach the full infective potential of
35 the virus.

 The use of mineral UV screens has also been described. Most do not absorb UV light but rather function by reflecting and refracting UV light. Major exceptions to this generalization are zinc oxide and titanium dioxide which have strong UV absorption

bands. Although titanium dioxide (TiO_2) is claimed in Japan [WPI Acc. #90-228659/30, *J. Econ. Entomol.* 69, 731(1976), and UK Patent Appl 2.043,448] to enhance virus stability toward UV radiation, it is also known that when exposed to ultraviolet light from sunlight it becomes reactive. For example, a sunlight irradiated aqueous suspension TiO_2 produced hydroxyl radicals which inactivated viruses in waste water treatment [*Water Research* 29 (1) 95, 1995].

It has been found that a modified form of TiO_2 , referred to as durable TiO_2 (as illustrated in U.S. Patent 4,125,412, EP 0 654 509 A1), provides a better and easier means of stabilization of baculoviruses than known stabilizers. In addition, other advantages of the use of durable TiO_2 will become apparent hereinafter.

SUMMARY OF THE INVENTION

The present invention pertains to a pesticidal composition comprising particles of biological pesticides and particles of durable TiO_2 , a modified form of TiO_2 , in an amount sufficient to substantially coat the particles of such biological pesticides; and an advantageous process for the preparation thereof. More specifically, the present composition pertains to an insecticidal baculovirus with an improved stabilizer.

The durable TiO_2 can be simply mixed with a biological pesticide, preferably though, durable TiO_2 is substantially coated on the surface of the biological pesticide. According to one preferred embodiment of the present invention, addition of an aqueous slurry of durable TiO_2 to a stirred aqueous slurry of a baculovirus at neutral pH (or alternately the virus to the TiO_2 slurry) causes the durable TiO_2 and the baculovirus to co-precipitate. Preferably, this material is then spray dried to particles less than 50 micron median volume diameter (mvd).

BRIEF DESCRIPTION OF THE PHOTOGRAPH

Figure 1 is an enlarged (40,000X) photo of the spray-dried product and shows a single coated virus.

DETAILED DESCRIPTION OF THE INVENTION

Microbial pesticides include the following organisms when they act as pesticides: 1) eucaryotic microorganisms including protozoa, algae and fungi; 2) procaryotic microorganisms including bacteria and; 3) viruses. Biological pesticides which are particularly enhanced by the present invention are those that are degraded by UV light, thus, quickly losing their efficacy. Such UV light degradable pesticides include, but are not limited to, baculoviruses, granulosis viruses, bacillus thuringensis, *Metarhizium* spp. and *Beauveria* spp.

One embodiment of this invention pertains to many different viruses belonging to more than a dozen families that have been isolated from insects. Among those, members of the family *Baculoviridae* are considered to hold most promise for commercial insect control. More detailed information concerning the various aspects of

baculoviruses as biopesticides can be found in the following publications: Ignoffo (1968), *Viruses-Living Insecticides, Current Topics in Microbiology and Immunology*, 42, 129-167; Arif and Jamieson (1989), *The Baculoviruses, Biocontrol of Plant Diseases, Vol. 1* (K.G. Mukerji & K.L. Garg, eds.); Blissard and Rohrmann (1990), *Baculovirus Diversity and Molecular Biology, Annual Review of Entomology* 35, 127-155; Adams and Bonami (1991), *Atlas of Invertebrate Viruses*, CRC Press, Boca Raton; Leisy and van Beek (1992), *Baculoviruses, Possible Alternatives to Chemical Insecticides, Chemistry & Industry*, April 1992, pp. 250-254; Vlak (1993), Genetic engineering of baculoviruses for insect control in: "*Molecular Approaches to Fundamental and Applied Entomology*" (Oakeshott and Whitten, eds.), Springer Verlag, New York; WO 96/36712.

As pertains to the present invention, the term baculovirus embraces insecticides containing a nuclear polyhedrosis virus or granulosis virus. As will be appreciated, in relation to microbial insecticides, in general the microbial pathogens themselves, known as occlusion bodies, are not in practice isolated from the mediums in which they are prepared and processed but can be used in a heterogeneous mixture containing a range of materials from the nutrients to the host's metabolic waste products. Particularly preferred embodiments of this invention embrace insecticides containing, but not limited to, the nuclear polyhedrosis viruses and granulosis viruses for the bollworm (*Helicoverpa zea*), the cotton bollworm (*H. armigera*), the tobacco budworm (*Heliothis virescens*), the cabbage looper (*Trichoplusia ni*), the fall armyworm (*Spodoptera frugiperda*), the beet armyworm (*S. exigua*), the cotton leafworm (*S. littoralis*), the pink bollworm (*Pectinophora gossypiella*), the Douglas-fir tussock moth (*Orgyia pseudotsugata*), the spruce budworm (*Choristoneura fumiferana*), the western spruce budworm (*C. occidentalis*), the gypsy moth (*Lymantria dispar*), the European pine sawfly (*Neodiprion sertifer*) the diamondback moth (*Plutella xylostella*), the grapeleaf skeletonizer (*Harrisina americana*), the potato tuber worm (*Phthorimeaea operculella*), the black cutworm (*Agrotis ipsilon*), the codling moth (*Cydia pomonella*), the oriental fruit moth (*C. molesta*), and the navel orangeworm (*Amyelois transitella*).

Baculovirus production is known to those skilled in the art and occurs by fermentation in cultured insect cells (Vlak et. al., 1990, *Bioreactor Development for Production of Viral Pesticides or Heterologous Proteins in "Insect Cell Cultures," Ann. N.Y. Acad. Sci.* 589, 399-418; Taticek et al., 1995, Overview of issues in bioreactor design and scale-up, in: "*Baculovirus Expression Systems and Biopesticides*," Shuler, Wood, Granados and Hammer, eds. Wiley-Liss, New York), or, *in vivo* in mass reared insect larvae (Ignoffo, 1967, Possibilities of Mass-Producing Insect Pathogens, *Internat. Colloq. Insect Pathol.* Netherlands, pp. 99-117; Shapiro 1986, *In Vivo Production of*

Baculoviruses in: *The Biology of Baculoviruses, Vol. II*, CRC Press; Shieh 1989, Industrial production of viral pesticides, *Adv. Virus Res.* 36, 315-343).

Durable TiO_2 means TiO_2 (either anatase or rutile) that has been made durable by coating the TiO_2 particles with an oxide coating consisting of alumina, silica, or a
5 combination of the two. An example of such a material is "TI-PURE®" Titanium Dioxide Pigment R-706 which is manufactured by the DuPont Company, Wilmington, DE. It is a TiO_2 on whose surface is deposited successive layers of silica and alumina which greatly reduces the photochemical reactivity of TiO_2 . Any
10 combination of silica or alumina or both may be contemplated however the preferred is one which has alumina as the outside layer. Preparation of these materials is described in *Inorganic Pigments, Manufacturing Processes*, Chemical Technology Review No. 166, M. H. Gutcho, ed. Noyes Data Corp. Park Ridge, NJ 1980 and references cited therein. The important feature is that the alumina, silica, etc., should form a continuous, nonporous coating on the TiO_2 particles and not be present as a simple admixture.
15 Typically these materials contain 1 to 4% silica (as SiO_2) and/or 4 to 9% alumina (as Al_2O_3), both based on the weight of TiO_2 .

The durable TiO_2 can be simply mixed with the virus at a weight ratio of 10 to 90% weight/weight (w/w) TiO_2 to virus. Preferably though, the durable TiO_2 is
20 substantially coated on the surface of the virus. Substantially coated means the TiO_2 is electrostatically fixed to the surface of the virus particle. Combination of the virus with the durable TiO_2 can occur by any standard means known in the art [*J. Econ. Entomol.* 69, 731(1976)] but is advantageously coated onto the individual virus particles by the process of the present invention. It is most preferred that the biological pesticide
25 particles be completely coated as individual particles or as heteroflocculated particles (i.e., some interior particles may not be coated or are partially coated, but the outside of the heteroflocculated particles are completely coated).

According to one preferred embodiment of the present invention, addition of an aqueous slurry of the R-706 durable TiO_2 particles to a stirred aqueous slurry of a
30 baculovirus at neutral pH (typically between about 5.5 and 8.0) causes the TiO_2 particles and the baculovirus solids to co-precipitate leaving a somewhat clear layer at the top of the liquid and a dense floc at the bottom. This is not observed with uncoated TiO_2 products (anatase or rutile) or TiO_2 products containing anionic surfaces such as silica-coated or those containing anionic surface-active agents. By virtue of its alumina
35 coating, the durable TiO_2 particles are positively charged in aqueous suspension at neutral pH whereas the nuclear polyhedrosis virus, and by inference the other impurities in the baculovirus, are negative. The existing charge difference and mutual attraction causes the system to "collapse" and produces the observed flocculation. This is a very beneficial effect in that it results in the TiO_2 being electrostatically (as opposed to

covalently) bound to the surface of the virus; thereby providing a very efficient means of surrounding each virus particle with a reversibly attachable UV shading material. This material is then spray dried to agglomerate the flocs into particles less than 50 micron mean volume diameter. This step provides resistance to redispersion of the durable TiO_2 away from the virus, which might occur during use in the presence of aggressive waters or surface active agents commonly employed in agricultural sprays, such as wetters or spreader-stickers while providing rapid dispersal under the very alkaline conditions of the target insect mid-gut where the pH can be ≥ 10.5 .

Certain particles which have light-filtering attributes but which bear a negative charge may be converted to bear a positive charge in which case the processes and benefits observed with durable TiO_2 are also observed. For example, a 5% suspension of Raven 430 Carbon (Columbian Chemicals Co.) containing 0.5% w/w of Ethoquad T/13-27W (Akzo Nobel Chemicals, Inc.) becomes positively charged. This suspension is also capable of flocculation like durable TiO_2 with the virus. Particles which have light filtering attributes include, but are not limited to TiO_2 , carbon and zinc oxide. Any means for converting the particles to bear a positive charge is acceptable, however, coating the particles as described for durable TiO_2 and use of suitable quaternary ammonium surfactants are preferred such as Ethoquad.

In commercial operation, the present composition will usually, but not necessarily, be diluted in an aqueous matrix and sprayed on the locus to be treated.

The present composition can optionally include inert ingredients typically found in agricultural formulations such as mineral carriers or diluents, surfactants to enhance wetting of the spray target, vegetable flours, animal proteins or other gustatory stimulants to encourage feeding on the formulations, buffering agents to maintain an optimum pH for the formulation, antifoam agents, and preserving or stabilizing agents to inhibit microbial growth or other degradative processes.

The present composition can also contain up to about 50% w/w certain fluorescent brighteners or stilbenes; or, alternatively, be added separately to an aqueous spray solution of the present composition. These materials are well known to provide enhanced activity of a range of baculoviruses to their hosts and are active at concentrations typically ranging from 0.01 to 1% by weight of the spray liquid but useful at higher concentrations. Suitable stilbene materials are described in U. S. Patent Nos. 5,124,149 and 5,246,936.

Preferred stilbene compounds are the analogs of 4,4'-diamino-2,2'-stilbene disulfonic acid, namely a Blancophor (available from Mobay Chemicals, Pittsburgh, PA) such as Blancophor BBH, Blancophor MBBH, Blancophor BHC, etc., a Calcifluor White (available from Sigma Chemical, St. Louis, MO) such as Calcifluor White, Calcifluor White M2R, Calcifluor White ABT, Calcifluor White LD, Calcifluor White

RWP etc., a Leucophor (available from Sandoz Chemicals Corp., Charlotte, NC) such as Lucophor BS, Lucophor BSB, Lucophor EKB, Lucophor PAB, etc.; a Phorwite (available from Mobay Chemicals, Pittsburgh, PA) such as Phorwite AR, Phorwite BBU, Phorwite BKL, Phorwite CL, Phorwite RKK, etc., Blancophor BBH, Calcifluor White M2R, and Phorwite AR are the most preferred stilbene compounds.

EXAMPLE 1

This example illustrates the general procedure for producing the flocculated durable TiO₂/virus mixtures. One hundred grams of durable TiO₂ particles (DuPont TI-PURE® Grade R-706) was bead-milled for 20 min with 100 g of deionized water. The pH was adjusted during milling to a range of 7.0 to 8.0 with dropwise addition of sodium hydroxide. The resultant particle size of this positively charged suspension was 0.8 microns mvd (mean volume diameter). This slurry was added with high speed mixing to 500 g of virus [*Autographa californica multicapsid nuclear polyhedrosis virus (MNPV)*] aqueous slurry (15%w/w suspended solids) obtained by infection of *Heliothis virescens* larvae. The slurry had been de-oiled by centrifugation. The virus solids typically contain 1×10^{10} to 1×10^{11} virus occlusion bodies OBs per gram, the virus used in this example contained 2.4×10^{10} (OBs) per gram dry weight. The pH of the resulting slurry was lowered to 6.5 by dropwise addition of hydrochloric acid and stirred briefly to permit completion of the process. The resultant mixture was spray dried in a pilot-scale spray dryer at an inlet temperature of 185°C and an outlet temperature of 60 to 65°C. A total of 100 g of durable TiO₂ coated virus formulation was recovered nominally containing 42% (1×10^{10} virus OBs per gram). The virus concentration cannot be directly measured in these systems because release from the matrix is only achieved at pH levels high enough to begin dissolving the virus particles. The particle size of the spray-dried material was 20.2 microns mvd and an example of a single coated virus is shown in Figure 1.

EXAMPLE 2

This example illustrates the general procedure for preparing a mixture as in Example 1 with a different virus and the addition of a fluorescent brightner. Fifty grams of durable TiO₂ particles (DuPont TI-PURE® Grade R-706) and 25.2 g of a stilbene brightener (Blankophor BBH) were bead-milled for 20 min with 75 g of deionized water. The pH was adjusted during milling to a range of 7.0 to 8.0 with dropwise addition of sodium hydroxide. The resultant particle size was 1.1 microns mvd. This slurry was added with high speed mixing to 547 g of virus (*Helicoverpa zea SNPV*) aqueous slurry (20%w/w suspended solids). The virus solids typically contain 1×10^{10} to 1×10^{11} virus OBs per gram, the virus used in this example contained 5.9×10^{10} OBs per gram dry weight. The pH of the resultant slurry was lowered to 6.5 by dropwise addition of hydrochloric acid. The resultant mixture was dried in the same spray dryer

as in Example 1 at an inlet temperature of 185°C and an outlet temperature of 60 to 65°C. The durable TiO₂ coated virus material was calculated to contain 5.3x10⁹ OBs per gram. The particle size of the spray-dried material was 18.9 microns mvd.

EXAMPLE 3

5 This example illustrates the general procedure for preparing a mixture as in Example 1 with a different virus and the addition of a feeding stimulant. Thirty-five grams of durable TiO₂ particles (DuPont TI-PURE® Grade R-706) and 65 g of corn flour were bead-milled for 20 min with 516 g of deionized water. The pH was adjusted during milling to a range of 7.0 to 8.0 with dropwise addition of sodium hydroxide. The
10 resultant particle size was 2.2 microns mvd. This slurry was added with high speed mixing to an aqueous slurry of 500 g of the virus of Example 1 (*Autographa californica MNPV*) slurry (25%w/w suspended solids). The virus solids typically contain 1x10¹⁰ to 1x10¹¹ OBs per gram, the virus used in this example contained 2.4x10¹⁰ OBs per gram dry weight. The pH of the resultant slurry was lowered to 6.5 by dropwise addition of
15 hydrochloric acid. The resultant mixture was dried in a spray dryer as in Example 1. The durable TiO₂ coated material was calculated to contain 1.3x10¹⁰ virus OBs per gram. The particle size of the spray-dried material was 26.8 microns mvd.

EXAMPLE 4

20 This example illustrates the general procedure for preparing a mixture as in Example 1 with a different virus and the addition of a binder. Twenty-five grams of durable TiO₂ particles (DuPont TI-PURE® Grade R-706) and 10 g of corn starch were bead-milled for 20 min with 75 g of deionized water. The pH was adjusted during milling to a range of 7.0 to 8.0 with dropwise addition of sodium hydroxide. This slurry was added with high speed mixing to 200 g of an aqueous slurry of virus (*Autographa californica MNPV*) slurry (20%w/w suspended solids). The virus solids typically
25 contain 1x10¹⁰ to 1x10¹¹ OBs per gram, the virus used in this example contained 2.4x10¹⁰ OBs per gram dry weight. The pH of the resultant slurry was lowered to 6.5 by dropwise addition of hydrochloric acid. The resultant mixture was spray dried in a spray dryer as in Example 1. The durable TiO₂ coated material was calculated to
30 contain 1.3x10¹⁰ virus OBs per gram.

EXAMPLE 5

35 This example illustrates the field stability achieved using the materials and process indicated in Example 1. Plots consisted of two 100 ft rows divided into four 25 ft replicates. Treatments were applied at 20 gallons-per-acre (0.18 m³ per hectare) with a 0.125%w/w aqueous suspension of unformulated virus (*Autographa californica MNPV*) using an application rate of 2.5x10¹¹ OBs per acre (1x10¹⁰ OBs per gram of virus). Another row of cotton was treated in the same manner and occlusion body rate using a 0.66% w/w aqueous dispersion of the powder described in Example 1. At

various time intervals whole cotton leaf samples were taken from each treated row along with an uncontaminated check row. Following application, plants were allowed to dry then 16 leaves per replicate was excised for laboratory bioassay. A single leaf was placed into each cell of a 16 celled jelly tray. A single 3-day-old *H. virescens* larva was also added to each cell. Test units were held at 28°C in 14 h of light for a total of 5 days. Mortality was assessed on the fifth day. Repeated leaf collections and bioassays were conducted 1, 2, and 4 d following the application. The results shown in Table 1 illustrate the stabilization achieved on addition of the durable TiO₂ particles.

10

Table 1
Comparison of formulated virus and unformulated virus

Sampling Time	Percent Kill	
	Unformulated	Formulation in Example 1
0	58	49
1 day	30	72
2 days	16	56
4 days	6	47

EXAMPLE 6

15 This example compares the activity of formulations prepared as in Examples 1 and 3 (with and without spray drying) as well as showing that the activity differences are maintained on a different surface (glass) as well as addition of a commonly used agricultural surfactant Tween 20®. In this Example, microscope slide coverslips were treated with 10 µl of virus preparation containing 1x10⁸ OBs per ml as calculated from the amount contained in each preparation and the suspension also contained 0.2% w/w Tween-20® surfactant. Half of the treated coverslips were exposed to natural mid-day sunlight for 4 h and the other half kept in darkness. The virus was stripped from each of the coverslips by sonication (10 s) in 10 ml of distilled water. A molten lepidopteran diet was added to the suspension of each sample and then cast into 2-25 well bioassay trays. Each replicate consisted of fifty 3-day-old

20 *H. virescens* larvae. Mortality assessments were made 144 h post exposure. The percent activity remaining was calculated by comparing UV exposed to UV unexposed virus. This example also illustrates the importance of spray drying the product prior to use.

30

Table 2
Comparison of formulations prepared according to Examples 1 and 3
after 4 hours of sun exposure.

Example 1	Example 3	Example 1 w/o Spray Drying
82	96	17

CLAIMS

What is claimed is:

1. A pesticidal composition comprising particles of biological pesticides and
5 particles of durable TiO_2 in an amount sufficient to substantially coat the particles of biological pesticides.
2. A pesticidal composition of Claim 1 wherein the biological pesticides are particles of a baculovirus.
3. A pesticidal composition of Claim 2 wherein the baculovirus contains genes
10 encoding an insecticidal protein.
4. A pesticide composition of Claim 2 wherein the particles of baculovirus substantially coated with durable TiO_2 have a mean particle size distribution of less than about 50 μm .
5. A pesticidal composition of Claim 1 and at least one of a surfactant, a solid
15 diluent or a liquid diluent.
6. A pesticidal composition of Claim 5 which also contains a fluorescent brightener.
7. A process for preparing the composition of Claim 1 comprising mixing durable TiO_2 particles with an aqueous slurry of biological pesticide particles.
- 20 8. A process of Claim 7 wherein the pH of the slurry is 5.5 to 8.
9. A process of Claim 7 wherein the resulting aqueous slurry is spray dried.
10. A method of treating agricultural pests comprising: applying to the pests or their environment a pesticidally effective amount of the pesticidal composition of Claim 1.

1/1




FIG. 1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/17393

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N63/00 A01N25/26 A01N25/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N C09C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE WPI Week 9030 Derwent Publications Ltd., London, GB; AN 90-228659 XP002051437 & JP 02 157 204 A (HOKKO CHEM. IND. CO. LTD.) cited in the application see abstract</p>	1-10
A	<p>--- WO 95 33378 A (LIGNOTECH USA INC) 14 December 1995 see page 1 - page 5, line 24 see page 8, line 12 - page 9, line 12 --- -/--</p>	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 January 1998

Date of mailing of the international search report

20/01/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Muellners, W

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/17393

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 573 150 A (TIOXIDE GROUP SERVICES LTD) 8 December 1993 see page 2, line 1 - line 48 see page 3, line 31 - line 57 ----	1-10
A	D.L. BULL ET AL.: "Improved Formulations of the Heliothis Nuclear Polyhedrosis Virus" JOURNAL OF ECONOMIC ENTOMOLOGY, vol. 69, no. 6, December 1976, pages 731-6, XP002051436 cited in the application see the whole document ----	1-10
A	US 4 115 144 A (CHAMBERS HUBERT HAROLD ET AL) 19 September 1978 see column 1 - column 3, line 19; claims ----	1-10
A	US 4 125 412 A (WEST WILLIAM A) 14 November 1978 cited in the application see column 1 - column 2, line 21 -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/17393

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9533378 A	14-12-95	US 5529772 A AU 2821595 A CA 2190655 A EP 0763974 A FI 964821 A	25-06-96 04-01-96 14-12-95 26-03-97 02-12-96
EP 0573150 A	08-12-93	AU 661773 B CA 2092239 A GB 2267503 A, B JP 6100701 A MX 9303330 A NO 931987 A NZ 247762 A US 5509960 A US 5554215 A ZA 9303347 A	03-08-95 05-12-93 08-12-93 12-04-94 29-07-94 06-12-93 27-06-95 23-04-96 10-09-96 17-11-93
US 4115144 A	19-09-78	GB 1560944 A AU 503276 B AU 2064176 A FR 2336169 A	13-02-80 30-08-79 22-06-78 22-07-77
US 4125412 A	14-11-78	AU 514147 B AU 2860677 A BR 7705970 A CA 1113658 A DE 2740561 A FR 2364252 A GB 1589070 A JP 1011670 B JP 1539263 C JP 53033228 A	29-01-81 15-03-79 27-06-78 08-12-81 16-03-78 07-04-78 07-05-81 27-02-89 16-01-90 29-03-78